



**Package leaflet: Information for the user**  
**SUNITINIB AQVIDA 12.5 mg hard capsules**  
**SUNITINIB AQVIDA 25 mg hard capsules**  
**SUNITINIB AQVIDA 37.5 mg hard capsules**  
**SUNITINIB AQVIDA 50 mg hard capsules**  
**Sunitinib**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet

1. What Sunitinib AqVida is and what it is used for
2. What you need to know before you take Sunitinib AqVida
3. How to take Sunitinib AqVida
4. Possible side effects
5. How to store Sunitinib AqVida
6. Contents of the pack and other information

## 1. What Sunitinib AqVida is and what it is used for

Sunitinib AqVida contains the active substance sunitinib, which is a protein kinase inhibitor. It is used to treat cancer by preventing the activity of a special group of proteins which are known to be involved in the growth and spread of cancer cells.

Sunitinib AqVida will only be prescribed to you by a doctor with experience in the use of anti-cancer medicinal products.

Sunitinib AqVida is used to treat adults with the following types of cancer:

- Gastrointestinal stromal tumour (GIST), a type of cancer of the stomach and bowel, where imatinib (another anticancer medicine) no longer works or you cannot take imatinib.
- Metastatic renal cell carcinoma (MRCC), a type of kidney cancer that has spread to other parts of the body.
- Pancreatic neuroendocrine tumours (pNET) (tumours of the hormone-producing cells in the pancreas) that have progressed or cannot be removed with surgery.

If you have any questions about how Sunitinib AqVida works or why this medicine has been prescribed for you, ask your doctor.

## 2. What you need to know before you take Sunitinib AqVida

### Do not take Sunitinib AqVida

- If you are allergic to sunitinib or any of the other ingredients of Sunitinib AqVida.

### Warnings and precautions

#### Talk to your doctor before taking Sunitinib AqVida:

**- If you have high blood pressure.** Sunitinib AqVida can raise blood pressure. Your doctor may check your blood pressure during treatment with Sunitinib AqVida, and you may be treated with medicines to reduce the blood pressure, if needed.

**- If you have or have had blood disease, bleeding problems, or bruising.** Treatment with Sunitinib AqVida may lead to a higher risk of bleeding or lead to changes in the number of certain cells in the blood which may lead to anaemia or affect the ability of your blood to clot. If you are taking warfarin or acenocoumarol, medicines which thin the blood to prevent blood clots, there may be a greater risk of bleeding. Tell your doctor if you have any bleeding while on treatment with Sunitinib AqVida.

**- If you have heart problems.** Sunitinib AqVida can cause heart problems. Tell your doctor if you feel very tired, are short of breath, or have swollen feet and ankles.

**- If you have abnormal heart rhythm changes.** Sunitinib AqVida can cause abnormality of your heart rhythm. Your doctor may obtain electrocardiograms to evaluate for these problems during your treatment with Sunitinib AqVida. Tell your doctor if you feel dizzy, faint, or have abnormal heartbeats while taking Sunitinib AqVida.

**- If you have had a recent problem with blood clots in your veins and/or arteries (types of blood vessels), including stroke, heart attack, embolism, or thrombosis.** Call your doctor immediately if you get symptoms such as chest pain or pressure, pain in your arms, back, neck or jaw, shortness of breath, numbness or weakness on one side of your body, trouble talking, headache, or dizziness while on treatment with Sunitinib AqVida.

**- If you have or have had damage to the smallest blood vessels known as Thrombotic Microangiopathy (TMA).** Tell your doctor if you develop fever, fatigue, tiredness, bruising, bleeding, swelling, confusion, vision loss and seizures.

**- If you have thyroid glands problems.** Sunitinib AqVida can cause thyroid gland problems. Tell your doctor if you get tired more easily, generally feel colder than other people or your voice deepens whilst taking Sunitinib AqVida. Your thyroid function should be checked before you take Sunitinib AqVida and regularly while you are taking it. If your thyroid gland is not producing enough thyroid hormone, you may be treated with thyroid hormone replacement.

**- If you have or have had pancreatic or gallbladder disorders.** Tell your doctor if you develop any of the following signs and symptoms: pain in the area of the stomach (upper abdomen), nausea, vomiting, and fever. These may be caused by inflammation of the pancreas or gallbladder.

**- If you have or have had liver problems.** Tell your doctor if you develop any of the following signs and symptoms of liver problems during Sunitinib AqVida treatment: itching, yellow eyes or skin, dark urine, and pain or discomfort in the right upper stomach area. Your doctor should do blood tests to check your liver function before and during treatment with Sunitinib AqVida, and as clinically indicated.

**- If you have or have had kidney problems.** Your doctor will monitor your kidney function.

**- If you are going to have surgery or if you had an operation recently.**

Sunitinib AqVida may affect the way your wounds heal. You will usually be taken off Sunitinib AqVida if you are having an operation. Your doctor will decide when to start Sunitinib AqVida again.

**- You may be advised to have a dental check-up before you start treatment with Sunitinib AqVida.**

- if you have or have had pain in the mouth, teeth and/or jaw, swelling or sores inside the mouth, numbness or a feeling of heaviness in the jaw, or loosening of a tooth tell your doctor and dentist immediately.

- if you need to undergo an invasive dental treatment or dental surgery, tell your dentist that you are being treated with Sunitinib AqVida in particular when you are also receiving or have received intravenous bisphosphonates. Bisphosphonates are medicines used to prevent bone complications that may have been given for another medical condition.

**- If you have or have had skin and subcutaneous tissue disorders.** While you are on this medicine "pyoderma gangrenosum" (painful skin ulceration) or "necrotising fasciitis" (rapidly spreading infection of the skin/soft tissue that may be life-threatening) may occur. This event is generally reversible after drug discontinuation. Severe skin rashes (Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme) have been reported with the use of sunitinib, appearing initially as reddish target-like spots or circular patches often with central blisters on the trunk. The rash may progress to widespread blistering or peeling of the skin and may be life-threatening. If you develop a rash or these skin symptoms, seek immediate advice from a doctor.

**- If you have or have had seizures.** Notify your doctor as soon as possible if you have high blood pressure, headache, loss of sight.

**- If you have diabetes.** Blood sugar levels in diabetic patients should be checked regularly in order to assess if anti-diabetic drug dosage needs to be adjusted to minimize the risk of low blood sugar.

### Children and adolescents

Sunitinib AqVida is not recommended for people aged under 18.

### Other medicines and Sunitinib AqVida

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription even those not prescribed.

Some medicines can affect the levels of Sunitinib AqVida in your body. You should inform your doctor if you are taking medicines containing the following active substances:

- ketoconazole, itraconazole – used to treat fungal infections
- erythromycin, clarithromycin, rifampicin – used to treat infections
- ritonavir – used to treat HIV
- dexamethasone – a corticosteroid used for various conditions
- phenytoin, carbamazepine, phenobarbital – used to treat epilepsy and other neurological conditions
- herbal preparations containing St. John's Wort (Hypericum perforatum) – used to treat depression and anxiety

### Sunitinib AqVida with food and drink

You should avoid drinking grapefruit juice while on treatment with Sunitinib AqVida.

### Pregnancy and breast-feeding

If you are pregnant or think you may be, tell your doctor.

Sunitinib AqVida is not to be used during pregnancy unless clearly necessary. Your doctor will discuss with you the potential risk of taking Sunitinib AqVida during pregnancy.

If you might get pregnant, you should use reliable method of contraception during treatment with Sunitinib AqVida.

If you are breast-feeding, tell your doctor. Do not breast-feed during treatment with Sunitinib AqVida.

### Driving and using machines

If you experience dizziness or you feel unusually tired, take special care when driving or using machines.

### 3. How to take Sunitinib AqVida

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Your doctor will prescribe a dose that is right for you, depending on the type of cancer to be treated. If you are being treated for GIST or MRCC, the usual dose is 50 mg once daily taken for 28 days (4 weeks), followed by 14 days (2 weeks) of rest (no medicine), in 6-week cycles. If you are being treated for pNET, the usual dose is 37.5 mg once daily without a rest period.

Your doctor will determine the appropriate dose you need to take as well as if and when you need to stop treatment with Sunitinib AqVida.

Sunitinib AqVida can be taken with or without food.

### If you take more Sunitinib AqVida than you should

If you have accidentally taken too many capsules, talk to your doctor straight away. You may require medical attention.

**If you forget to take Sunitinib AqVida** Do not take a double dose to make up for a forgotten dose.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You must immediately contact your doctor if you experience any of those serious side effects (see also **What you need to know before you take Sunitinib AqVida**):

**Heart problems.** Tell your doctor if you feel very tired, are short of breath, or have swollen feet and ankles. These may be symptoms of heart problems that may include heart failure and heart muscle problems (cardiomyopathy).

**Lung or breathing problems.** Tell your doctor if you develop cough, chest pain, sudden onset of shortness of breath or coughing up blood. These may be symptoms of a condition called pulmonary embolism that occurs when blood clots travel to your lungs.

**Kidney disorders.** Tell your doctor if you experience altered frequency or absence of urination which may be symptoms of kidney failure.

**Bleeding.** Tell your doctor right away if you have any of these symptoms or a serious bleeding problem during treatment with Sunitinib AqVida: painful, swollen stomach (abdomen), vomiting blood, black, sticky stools, bloody urine, headache or change in your mental status, coughing up of blood or bloody sputum from the lungs or airway.

**Tumour destruction leading to hole in the intestine.** Tell your doctor if you have severe abdominal pain, fever, nausea, vomiting, blood in your stool, or changes in bowel habits.

Other side effects with Sunitinib AqVida may include:

*Very common side effects (may affect more than 1 user in 10)*

- Reduction in the number of platelets, red blood cells and/or white blood cells (e.g. neutrophils).
- Shortness of breath.
- High blood pressure.
- Extreme tiredness, loss of strength.
- Swelling caused by fluid under the skin and around the eye, deep allergic rash.
- Mouth pain/irritation, mouth sores/inflammation/dryness, taste disturbances, upset stomach, nausea, vomiting, diarrhoea, constipation, abdominal pain/swelling, loss/decrease of appetite.
- Decreased activity of thyroid gland (hypothyroidism).
- Dizziness.
- Headache.
- Nose bleeding.
- Back pain, joint pain.
- Pain in arms and legs.
- Yellow skin/skin discoloration, excess pigmentation of the skin, hair colour change, rash on the palms of the hands and soles of the feet, rash, dryness of the skin.
- Cough.
- Fever.
- Difficulty in falling asleep.

*Common side effects (may affect 1 to 10 users in 100)*

- Blood clots in the blood vessels.
- Deficiency of blood supply to the heart muscle, due to obstruction or constriction of the coronary arteries.
- Chest pain.
- Decreased in the amount of blood pumped by the heart.
- Fluid retention including around the lungs.
- Infections.
- Decreased blood sugar level. If you experience any signs and symptoms of low blood sugar: Notify your doctor as soon as possible if you experience fatigue, palpitations, sweating, hunger and loss of consciousness.
- Loss of protein in the urine sometime resulting in swelling.
- Influenza-like syndrome.
- Abnormal blood tests including pancreatic and liver enzymes.
- High level of uric acid in the blood.
- Haemorrhoids, pain in the rectum, gingival bleeding, difficulty in swallowing or inability to swallow.
- Burning or painful sensation in the tongue, inflammation of the digestive tract lining, excessive gas in the stomach or intestine.
- Weight loss.
- Musculoskeletal pain (pain in muscles and bones), muscular weakness, muscular fatigue, muscle pain, muscle spasms.
- Nasal dryness, congested nose.
- Excessive tear flow.
- Abnormal sensation of the skin, dry skin, itching, flaking and inflammation of the skin, blisters, acne, nail discoloration, hair loss.
- Abnormal sensations in extremities.
- Abnormally decreased/increased sensitivity, particularly to touch.
- Acid heartburn.
- Dehydration.
- Hot flushes.
- Abnormally coloured urine.
- Depression.
- Chills.

*Uncommon side effects (may affect 1 to 10 users in 1,000)*

- Life-threatening infection of the soft tissue including the ano-genital region. Contact your doctor immediately if symptoms of infection occur around a skin injury, including fever, pain, redness, swelling, or drainage of pus or blood.
- Stroke.
- Heart attack caused by an interrupted or decreased blood supply to the heart.
- Changes in the electrical activity or abnormal rhythm of the heart.
- Fluid around the heart (pericardial effusion).
- Liver failure.
- Pain in the stomach (abdomen) caused by inflammation of the pancreas.
- Tumour destruction leading to hole in the intestine (perforation).
- Inflammation (swelling and redness) of the gallbladder with or without associated gallstones.
- Abnormal tube like passage from one normal body cavity to another body cavity or the skin.

- Pain in the mouth, teeth and/or jaw, swelling or sores inside the mouth, numbness or a feeling of heaviness in the jaw, or loosening of a tooth. These could be signs and symptoms of bone damage in the jaw (osteonecrosis). Tell your doctor and dentist immediately if you experience any of them.
- Overproduction of thyroid hormones which increases the amount of energy the body uses at rest.

- Problems with wound healing after surgery.
- Increased blood level of enzyme (creatine phosphokinase) from muscle.
- Inappropriate and excessive reaction to an allergen.

*Rare side effect (may affect 1 to 10 users in 10,000)*

- Severe reaction of the skin and/or mucous membranes (Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme).
- Tumour lysis syndrome (TLS) – TLS consists of a group of metabolic complications that can occur during treatment of cancer. These complications are caused by the break-down products of dying cancer cells and may include the following: nausea, shortness of breath, irregular heartbeat, muscular cramps, seizure, clouding of urine and tiredness associated with abnormal laboratory test results (high potassium, uric acid and phosphorous levels and low calcium levels in the blood) that can lead to changes in kidney function and acute renal failure.
- Abnormal muscle breakdown which can lead to kidney problems (rhabdomyolysis).
- Abnormal changes in the brain that can cause a collection of symptoms including headache, confusion, seizures, and vision loss (reversible posterior leukoencephalopathy syndrome).
- Painful skin ulceration (pyoderma gangrenosum).
- Inflammation of the liver (hepatitis).
- Inflammation of the thyroid gland.
- Damage to the smallest blood vessels known as thrombotic microangiopathy (TMA).

### Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below).

Rooyan darou Company  
 No.20, Salman-e-pak Alley, Ostad Nejatollahi St., Post code: 15996-73617  
 Tehran-Iran

### 5. How to store Sunitinib AqVida

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the outer pack and label after "EXP". The expiry date refers to the last day of that month.
- This medicine does not require any special storage conditions.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### 6. Contents of the pack and other information

#### What Sunitinib AqVida contains

#### Sunitinib AqVida 12.5 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 12.5 mg sunitinib.

- *Capsule content:* mannitol, croscarmellose sodium, povidone (K-25) and magnesium stearate.

#### Sunitinib AqVida 25 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 25 mg.

- *Capsule content:* mannitol, croscarmellose sodium, povidone (K-25) and magnesium stearate.

#### Sunitinib AqVida 37.5 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 37.5 mg.

- *Capsule content:* mannitol, croscarmellose sodium, povidone (K-25) and magnesium stearate.

#### Sunitinib AqVida 50 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 50 mg.

- *Capsule content:* mannitol, croscarmellose sodium, povidone (K-25) and magnesium stearate.

**Marketing Authorization Holder:** Rooyan Darou pharmaceutical company  
**Under licence of AqVida - Germany**  
 No.13, Maragheh Alley, Ostad Nejatollahi St.  
 15996-77814 Tehran-IRAN

**This leaflet was last revised in june ,2016**